

BioLure® Fruit Fly

Registration No. L7695 Act No. 36 of 1947

A controlled release fruit fly attractant for use as a monitoring system with McPhail type traps on citrus, grapes, deciduous and subtropical fruit orchards against *Ceratitis spp.*

Active ingredients:

Ammonium Acetate	211 g/kg
Trimethylamine hydrochloride	91 g/kg
1,4-diaminobutane (Putrecine)	3 g/kg

Distributed by:

Chempac (Pty) Ltd
Reg. No 2002/007885/07
P.O. Box 516, Suider Paarl, 7624
Tel No. (021) 874 1055

Registered in RSA by:

Spectrum Research Services cc
Reg. No. 1997/59453/23
Suite 202, Postnet X2600
HOUGHTON, 2041

Green band



CAUTION
VERSIGTIG



WARNINGS:

- May cause skin and eye irritation.
- Store unused product in a refrigerator away from food.
- Keep out of reach of children, uninformed persons and animals.
- ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

- Avoid contact with skin, eyes and clothing. In case of accidental eye contact flush with plenty of clean water.
- Wash with soap and water after handling the dispensers.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.
- Destroy empty container and do not use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Method of application: Remove the protective strip from each of the three dispensers before placing them into the trap. The Mc Phail type trap is recommended for use with Biolure® Fruit Fly Lure. Insert a Vapona Strip tablet (Reg.no. L2187) into the bottom of the trap prior to placing in the orchard or vineyard. Hang the trap at head height ($\pm 1,8$ m) on an outside limb of a tree or vine (as high up in bush vines as possible).

Place one trap per 2 hectares of orchard / vineyard.

Ensure the trap entrance is free from foliage in order to allow free air movement and easy access for flies.

Replace the vapour dispensers and Vapona strip every 42 days. Remove all old replaced dispensers and protective strips from orchards.

Ensure that the trap is hung away from the working row to avoid damage from tractors, spray machines or direct hits from irrigation sprinklers

The interpretation of the results should be discussed with a local adviser.

NOTES:

5. It is important that the mouth of the trap be kept open and free from obstacles (twigs, leaves).