

# CUPPROFIX DISPERSS™

Registrasie Nr. L6784

Wet Nr. 36 van 1947

Registration No. L 6784

Act 36 of 1947

## BORDEAUX MENGSEL / MIXTURE

'n Water-dispergeerbare korrel swamdoder vir die beheer van sekere swamsiektes soos aangedui.

A water dispersible granular fungicide for the control of certain fungal diseases as listed.

### AKTIEWE BESTANDELE:

### ACTIVE INGREDIENTS:

Kopersulfaat  
Kalk  
(Kopersulfaat met kalk geneutraliseer)

770 g/kg  
174 g/kg

Copper sulphate  
Lime  
(Copper sulphate neutralised by sodium lime)

Metalliese koper

200 g /kg

Metallic copper

Netto Massa

**20 kg**

Net Mass

Vervaardig deur:  
Cerexagri S.A. 1 rue des Frères Lumière,  
BP 9, 78373 PLAISIR Cedex – Frankryk

Manufactured by:  
Cerexagri S.A. 1 rue des Frères Lumière,  
BP 9, 78373 PLAISIR Cedex - France

Geregistreer en Versprei deur:  
Chempac (Pty) Ltd., Co. Reg. No. 2002/007885/07  
Posbus 516, Suider-Paarl, 7624  
Tel. Nr. (021) 874 1055

 **chempac**  
Progressive agricoore

Registered and Distributed by:  
Chempac (Pty) Ltd., Co. Reg. No. 2002/007885/07  
P O Box 516, Suider Paarl, 7624  
Tel. No. (021) 874 1055

Lot Nr.


Batch No.

Datum van Vervaardiging

Date of Manufacture



**CAUTION**  
**VERSIGTIG**



**WAARSKUWINGS:**

Laat 14 dae toe tussen die laaste toediening en oes van appelkose, perkes en pruiwe, 21 dae vir wyndruive en 4-6 weke vir tafeldruive. Giftig indien ingesluk. Kan matige oog- en velirritasie veroorsaak. Giftig vir vis. Hou buite bereik van kinders, oningeligte persone en diere. Berg weg van voedsel en voer.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaat – en bergings toestande, kwaliteit van verduunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die patogene teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effekte op mens of dier of vir 'n gebrek aan prestasie as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

**VOORSORGMATREEL'S:**

Vermyn naseming van poeier of spuitreuel.  
Dra rubberhandskoene en 'n gesigskerm wanneer spuitmengsel voorberei word.  
Was na gebruik.  
Vermyn die oorvaai van spuitreuel na ander gewasse, riviere en damme.  
Vermyn besoedeling van voedsel, voer, eetgerei en drinkwater.  
Houers en verpakings moet heeltemal leeggemaak word voordat hulle vernietig word.  
Skud die houer deeglik uit in die mengbak of spuitenk voordat leë houer vernietig word en moenie vir enige ander doel gebruik nie.  
Maak spuitapparaat skoon na gebruik en raak ontslae van waswater waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.

**GEBRUIKSAANWYSING: Gebruik alleenlik soos aangedui.**

Meng die aanbevole hoeveelheid met water en roer deeglik. Bespuit die plante deeglik aan weerskante van die blare. CUPROFIX DISPERSS is 'n beskermende spuitmiddel en moet verkleeslik toegedien word voordat die siekte verskyn.

Gewas / Siekte	Dosis / 100 l water	Opmerkings
<b>Appelkose, Perskes</b>		
<b>Pruiwe:</b>	0,8 – 1,0 kg	1 <sup>ste</sup> bespuiting: nadat die vrugte geoes is. 2 <sup>de</sup> bespuiting: 3 – 4 weke later 3 <sup>de</sup> bespuiting: net voor knopswel
<b>Gonksiesiekte</b>		
<b>Bakteriese kanke</b>		
<b>Druive:</b>		
<b>Donskimmel</b>	1 kg	1 <sup>ste</sup> bespuiting: wanneer lote 2,5 – 5 cm lank is. Herhaal met nog 2 – 3 bespuitings met tussenposes van 14 dae
<b>Bakteriese skroesiekte,</b>		
<b>Antraknose en</b>		
<b>Streepvleksiekte</b>		

**WARNINGS:**

Allow 14 days between last application and harvest of apricots, peaches and plums, 21 days for wine grapes and 4-6 weeks for table grapes. Poisonous if swallowed. May cause mild eye and skin irritation. Toxic to fish. Keep out of reach of children, unformed persons and animals. Store away from food and feed.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climate and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pathogen against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

**PRECAUTIONS:**

Avoid inhalation of dust or spray mist.  
Wear rubber gloves and face shield when preparing spray mixtures.  
Wash after use.  
Avoid drift onto other crops, grazing, rivers and dams.  
Avoid contamination of food, feed, eating utensils and drinking water.  
Containers and packages must be completely emptied before being disposed of.  
Shake out thoroughly into the mixing or spray tank before destroying the empty container and do not use for any other purpose.  
Clean application equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

**DIRECTIONS FOR USE: use only as directed:**

Mix the recommended quantity with water and stir well. Spray the plants thoroughly on both sides of the leaves. CUPROFIX DISPERSS is a protective spray and should preferably be applied before disease appears.

Crop / Disease	Dosage / 100 l water	Remarks
<b>Apricots, Peaches</b>		
<b>Plums:</b>	0,8 – 1,0 kg	1 <sup>st</sup> spray: after fruit has been harvested. 2 <sup>nd</sup> spray: 3 – 4 weeks later 3 <sup>rd</sup> spray: just before bud movement.
<b>Gum spot and</b>		
<b>Bacterial canker</b>		
<b>Grapes:</b>		
<b>Downy mildew</b>	1 kg	1 <sup>st</sup> spray: when shoots are 2,5 – 5 cm long. Repeat with 2 – 3 further sprays at 14 days intervals.
<b>Bacterial blight,</b>		
<b>Anthraxnose and</b>		
<b>Dead arm</b>		