

CHEMPAC SODIUM MOLYBDATE

Sodium Molybdate Natrium Molibdaat

Reg. No./Nr. B 3457 : Act/Wet 36 of/van 1947

Fertiliser - Grp 2 - Misstof

Molybdenum (Mo) **395 g/kg** Molibdeen (Mo)

Mass 1 or 25 kg Massa

Registered by / Geregistreer deur: Chempac (Pty) Ltd Reg.No. / Nr. 2002/007885/07
P O Box / Posbus 516, Suider Paarl 7624.
Tel.No. Nr. (021) 874 1055



WARNING: Always consult a qualified agronomist before applying Chempac Sodium Molybdate

DIRECTIONS FOR USE: Use only as directed

CROP	DOSAGE	REMARKS
Maize	50 g / 1,25 ℓ	Place 100 kg seed in a drum with lid. Add the solution and rotate the drum for 15 to 20 minutes. The seed may be treated any time before planting.
Sunflowers	25 g / 25 kg seed 5g /100ℓ to a maximum of 75g/ha	Apply solution uniformly to the seed. Apply to seedlings as a full cover foliar spray.
Soya beans	75 – 125 g / ha	Apply to seedlings as a foliar spray.
Tomatoes	250 g / ha	Apply to planting furrow before transplanting or as a foliar spray.
Lucerne	125 – 250 g / ha	Seed treatment simultaneously with inoculation.
Cucurbits	2 g / ℓ	Let the seed soak overnight in a 0,2 % solution and plant immediately afterwards.
Crucifers	100 – 250 g / ha	Apply as a foliar spray to young plants using sufficient water.
Grass and Clover pastures	155 g / ha	Foliar spray onto young re-growth after cutting.

Depending on the amount of Chempac Sodium Molybdate applied, the following reapplication intervals are recommended:

< 75 g - 1 year
75 to 150g - 2 to 3 years
> 150 g - 4 to 5 years